



SPRING IS HERE.
IS YOUR DOG PROTECTED FROM LEPTOSPIROSIS?

LEPTOSPIROSIS

Leptospirosis is a dangerous bacterial infection
ALL DOGS ARE AT RISK OF CONTRACTING



LEPTOSPIROSIS
IN CATS IS RARE

RISK FACTORS
include exposure to:



wildlife



the urine of
an infected
pet or animal



contaminated
water or soil



Infected pets develop flu-like symptoms
such as fever, loss of appetite, vomiting, diarrhea,
dehydration, weakness, depression, and lethargy.

In serious cases, infection can be fatal. High concentrations
of bacteria in the liver and kidneys can cause organ failure.




LEPTOSPIROSIS IS
ZOONOTIC




(VACCINATION PROTECTS YOUR HUMAN FAMILY TOO!)

VACCINATION IS MORE AFFORDABLE THAN TREATMENT
AND WILL SPARE YOUR DOG FROM LIFE-THREATENING ILLNESS AND A PAINFUL RECOVERY.



HELP PROTECT YOUR DOG
FROM LYME DISEASE AND
TICK-BORNE INFECTIONS.

LYME DISEASE



LYME DISEASE IS A BACTERIAL,
TICK-TRANSMITTED DISEASE

Lyme disease is the fastest-growing infectious disease
and most common tick-borne disease in the country

WE FIND TICKS ON DOGS YEAR ROUND



Ticks are most prevalent in
WOODED, GRASSY AREAS

Inspect the skin + coat after spending time outdoors

COMMON SIGNS



fever



lethargy



painful joints;
limping



loss of appetite



kidney failure

Pets often show no clinical signs
until 2-5 months post-infection

ASK US ABOUT THE LYME VACCINE, ANNUAL TESTING, AND PREVENTATIVE RECOMMENDATIONS



CANINE INFLUENZA

CANINE INFLUENZA VIRUS IS A **HIGHLY CONTAGIOUS RESPIRATORY INFECTION**

THE VACCINE IS STRONGLY RECOMMENDED IF YOUR DOG...


-  travels regularly
-  stays at kennels
-  sees a groomer
-  goes to dog parks
-  attends daycare
-  enters dog shows

Causes flu-like symptoms similar to kennel cough, such as coughing, sneezing, lethargy, nasal discharge & runny eyes

In severe cases, pets can develop pneumonia, high fever, and difficulty breathing



NEARLY ALL DOGS EXPOSED TO THE VIRUS WILL CONTRACT IT



ABOUT 20% develop no clinical signs, but can spread infection

ABOUT 80% show clinical signs

Dogs that spend a lot of time with other dogs have the highest risk of exposure




THE PROTOCOL FOR VACCINATION CONSISTS OF AN INITIAL VACCINE, FOLLOWED BY A BOOSTER 3 WEEKS LATER, THEN ANNUALLY AS NEEDED




MICROCHIPS

TEN MILLION

pets are lost or stolen annually

Microchips **increase the chances** of a happy reunion 



1 in 3 pets will get lost in their lifetime

To date, HomeAgain Microchips has helped to return over **2 MILLION** pets to their families

Microchipped dogs are more than twice as likely to return home



Microchipped cats are more than twenty times as likely to return home



YOUR PET IS A PRICELESS ADDITION TO YOUR FAMILY.
WE WANT TO HELP MAKE SURE THEY STAY SAFE WITH YOU.